REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

1. REPORT DATE *(DD-MM-YYYY)*	2. REPORT TYPE	3. DATES COVERED *(From - To)*
	Technical Papers	

4. TITLE AND SUBTITLE

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

5d. PROJECT NUMBER
2303

5e. TASK NUMBER
M2C8

5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

8. PERFORMING ORGANIZATION REPORT

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S ACRONYM(S)

11. SPONSOR/MONITOR'S NUMBER(S)

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

11/21 013

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Leilani Richardson
Unclassified	Unclassified	Unclassified	A		19b. TELEPHONE NUMBER *(include area code)* (661) 275-5015

62 separate items are enclosed

Ann C8

MEMORANDUM FOR PR (In-House Publication)

FROM: PROI (TI) (STINFO) 30 Jun 2000

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2000-144**
M. Fajardo, S. Tam, "High Resolution Infrared Absorption Spectroscopy in Doped Parahydrogen Solids: CO/pH_2 – a Molecular Thermometer"

3rd International Conference on Cryocrystals and Quantum Crystals (Statement A)
(Szklarska Poreba, Poland, 28 Jul – 04 Aug 00) (Submission Deadline: 28 Jul 00)

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.
Comments:_____

Signature _____ Date _____

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review.
Comments: _____

Signature _____ Date _____

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b.) appropriateness of distribution statement, c.) military/national critical technology, d.) economic sensitivity, e.) parallel review completed if required, and f.) format and completion of meeting clearance form if required
Comments:_____

Signature _____ Date _____

4. This request has been reviewed by PR for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/ national critical technology, and f.) data rights and patentability
Comments:_____

APPROVED/APPROVED AS AMENDED/DISAPPROVED

_____

LESLIE S. PERKINS, Ph.D (Date)
Staff Scientist
Propulsion Directorate

3rd International Conference on Cryocrystals and Quantum Crystals, 28 Jul - 4 Aug 2000, Szklarska Poreba, Poland

High Resolution Infrared Absorption Spectroscopy in Doped Parahydrogen Solids: CO/pH_2 -- a Molecular Thermometer

Mario E. Fajardo, and Simon Tam
USAF Research Laboratory, AFRL/PRSP, Bldg. 8451, Edwards AFB, CA 93524-7680
mario_fajardo@plc.af.mil

* HEDM Cryosolid Propellants
* Trapping of Metal Atoms in Cryogenic Solid Hydrogen
* Rapid Vapor Deposition of Transparent Parahydrogen (pH_2) Solids
* High Resolution IR Absorption Spectroscopy in Doped pH_2 Solids
* CO/pH_2 "Thermometer" Depositions
* Summary

HEDM Cryosolid Propellants Payoffs

Increased Specific Impulse

$$I_{sp} \propto \sqrt{\Delta H_{sp}}$$

$LOX/LH_2 : I_{sp} = 390$ s

$5\%\ B/sH_2 + LOX : I_{sp} = 500$ s (+30%)*

* calculated for $P_{chamber} = 1000$ PSIA, $P_{exhaust} = 14.7$ PSIA

Greater Propellant Density

liquid H_2 @ 20 K : $\rho = 0.070$ g/cm^3

solid H_2 @ 2 K : $\rho = 0.087$ g/cm^3 (+25%)

50/50 liquid He/solid H_2 : $\rho = 0.105$ g/cm^3 (+50%)

Dopant recombination/reaction in solid pH_2

* ideally:

$$X + pH_2 \xrightarrow{T=2K} X/pH_2 \qquad \text{isolated atoms}$$

* in practice:

$$X + X + M \rightarrow X_2 + M \qquad \text{recombination}$$
$$\rightarrow X_n$$

$$X + H_2 + M \rightarrow HX + H + M \qquad \text{reaction}$$
$$\rightarrow H_nX + M$$

$$X_n + H_2 + M \rightarrow HX_n + H + M \qquad \text{both}$$
$$\rightarrow H_mX_n + M$$

Experimental Diagram (c1993)



M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

Transmission Spectrum of Li/nH_2, $d \approx 10\ \mu$



—— difference + 1.0

—·— pure pH_2

------ Li/pH_2

M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

Optical Scattering in Solid Hydrogen

Crystal Growing and Quality (p. 81)

"There is a considerable art to growing hydrogen crystals of high quality. Good crystals are always grown slowly from the melt; a rapid freeze from the gas produces snow."

Crystallite Light Scattering (p. 83)

"The reason that a good hydrogen crystal is so hard to see is its low refractive index...an estimated 1.16!

Yet a 1 mm-thick layer of hydrogen crystallites can be a completely opaque brown-black."

[P.C. Souers, Hydrogen Properties for Fusion Energy (UC Press, Berkeley, 1986)]

ortho- and para-hydrogen



[I.F. Silvera, Rev. Mod. Phys. **52**, 393 (1980)]

Experimental Diagram (c1997)



M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).
S. Tam and M.E. Fajardo, Rev. Sci. Instrum. **70**, 1926 (1999).

Transmission Spectrum of B/pH_2 d≈1mm



M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).
S. Tam, M. Macler, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys., submitted (2000).

IR Absorption of 6 mm Thick pH_2 Solid



Non-observation of the $Q_1(0)$ transition demonstrates the absence of oH_2 impurities, and that the microscopic structure is not amorphous or porous.

Observation of $S_1(0)$ transition demonstrates the absence of inversion symmetry for <u>some</u> H_2 molecular environments.

[van Kranendonk and Gush, Phys. Lett. **1**, 22 (1962)]

M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).

Raman Spectra of pH_2 Solids



Mixed hcp/fcc as-deposited structure, anneals to hcp.
[G.W. Collins, et al., Phys. Rev. B **53**, 102 (1996)]

(d) sample in (c) warmed to 4.5 K.
(c) 4.5 mm sample as deposited at 3.3 K (Φ = 290 mmol/hr).

(b) sample in (a) warmed to 4.5 K.
(a) 6 mm sample as deposited at 3.1 K (Φ = 200 mmol/hr).

M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).

High Res. IR Spectroscopy in Solid pH_2



FIG. 1. Apparatus for the simultaneous spectroscopy of the infrared and Raman transitions. The nonlinearity of $LiNbO_3$ is used for the former and that of solid H_2 is used for the latter. D.M., dichroic mirror; S. A., spectrum analyzer; P. B., polarizer beamsplitter.

[T. Momose, K.E. Kerr, D.P. Weliky, C.M. Gabrys, R.M. Dickson and T. Oka, J. Chem. Phys. **100**, 7840 (1994)]

ν_4 CH_4/pH_2 IR Absorptions (res = 0.01 cm^{-1})



S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

ν_4 CH_4/pH_2 IR Absorptions



(a) Rapid Vapor Deposited sample: as-deposited at 2.4 K
(b) Rapid Vapor Deposited sample: annealed to 4.8 K
(c) Enclosed Cell Condensed sample: cooled to 4.8 K

S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

ν_4 CH_4/pH_2 Energy Levels



S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

CO_2/pH_2 Trapping Sites



S. Tam and M.E. Fajardo, Fiz. Nizk. Temp. [Low Temp. Phys.] accepted (2000).

CO_2/pH_2 IR Absorptions (res = 0.008 cm^{-1})



as-deposited at T = 2.4 K

(c) 1.2 ppm CO_2/pH_2

(b) 0.04 ppm CO_2/pH_2

(a) 0.01 ppm CO_2/pH_2

S. Tam and M.E. Fajardo, Fiz. Nizk. Temp. [Low Temp. Phys.] accepted (2000).

$^{13}C^{16}O_2/pH_2$ vs. $^{12}C^{16}O_2/pH_2$



annealed, T = 2.4 K

d = 1.0 mm

2 ppm $^{13}C^{16}O_2$ and
2 ppm $^{12}C^{16}O_2$

88 PPM HCl/pH_2



Gas Phase $(HCl)_2$

High resolution, jet-cooled infrared spectroscopy of $(HCl)_2$: Analysis of ν_1 and ν_2 HCl stretching fundamentals, interconversion tunneling, and mode-specific predissociation lifetimes

Michael D. Schuder,[a] Christopher M. Lovejoy,[b] Robert Lascola,[c] and David J. Nesbitt[d]
Joint Institute for Laboratory Astrophysics, National Institute of Standards and Technology and University of Colorado, and the Department of Chemistry and Biochemistry, University of Colorado, Boulder, Colorado 80309

(Received 5 April 1993; accepted 7 June 1993)



FIG. 1. Vibrationally averaged structure and internal coordinates for HCl dimer. The intermolecular axis R connects the HCl centers of mass. The internal angles, θ_1 and θ_2, are measured from the intermolecular axis to the HCl bonds r. The torsion angle, $\phi=\phi_1-\phi_2$, is shown at 180° (planar). The minimum energy configuration shown is for $\theta_1=16°$, $\theta_2=87°$ with $\phi_1-\phi_2=180°$. The HCl subunit on the left is referred to as the bonded HCl with an associated vibration labeled ν_2. The proton on the other HCl is not involved with the hydrogen bond, and this subunit is referred to as the free HCl, with a vibration labeled ν_1.



J. Chem. Phys. v99, p4346 (1993).

$(HCl)_2/pH_2$ isotopomers



analysis in collaboration with D.T. Anderson, U. Wyoming.

Co-operative IR absorptions



analysis in collaboration with R.J. Hinde, U. Tennessee, Knoxville.

80 PPM CO/pH_2 (res = 0.1 cm^{-1})



(a) as deposited at T = 2.4 K

(b) warmed to T = 4.8 K

(c) re-cooled to T = 2.4 K

analysis in collaboration with T. Momose, Kyoto U.

$(CO)_n/pH_2$ IR Absorptions



≈ 6 μmol CO in each sample

d = 1.7 mm

d = 20 μm

d = 4.7 μm

Experimental Diagram



Top View

Side View

Experimental Protocol



Thermometer experiments:



control experiment

CO_2/pH_2 Thermometer Peak





Calibration of CO/pH_2 Thermometer



simplified Boltzmann:

$$A_{2135} = N \exp(-E/T_{CO})$$

$N = 0.0860$
$E = 7.896$ K

include $Q_{rot} \Rightarrow E \approx 11$ K

$6\ B_{CO/pH2} \approx 12$ K

Deposition Temperatures



#3:

Φ_{H2} = 240 mmol/hr

$\dot{R}$ = 55 μm/min

#2:

Φ_{H2} = 200 mmol/hr

$\dot{R}$ = 48 μm/min

#1:

Φ_{H2} = 110 mmol/hr

$\dot{R}$ = 26 μm/min

Cryostat Thermal Response



T_B: slope = 3.6 K/W

T_A: slope = 1.0 K/W

Deposition Heat Loads



heat fluxes calculated from
$\Delta T = T_B - T_B$(prior dep.)

slope of fit line $\Rightarrow$
$E_{dep} = 3.3$ kJ/mol

compare with:
$E_{vap} = 1.1$ kJ/mol

$\leftarrow$ note post-deposition radiative(?) heat loads

3rd International Conference on Cryocrystals and Quantum Crystals, 28 Jul - 4 Aug 2000, Szklarska Poreba, Poland

pH_2 Thermal Conductivity Calculations

expt.#	Δx(cm)	ΔT_{max}(K)	ΔT_{min}(K)	$(\dot{Q}/A)_{max}$	$(\dot{Q}/A)_{min}$	κ_{min}	κ_{max}
1	0.108	0.61	0.21	9.2	6.9	1.2	4.7
2	0.203	1.19	0.79	13.8	10.7	1.8	3.5
3	0.232	1.74	1.34	16.6	12.5	1.7	2.9

Units: $\dot{Q}/A$ (mW/cm^2), κ (mW/cm-K).

Summary: κ = **3(±2) mW/cm-K**, averaged over $2 < T < 5$ K range.

Comparison with Literature



FIG. 1. Thermal conductivity of crystals of pure pH_2 and pH_2 with Ne impurity (the concentration in ppm are indicated); the solid lines are calculated results.

Previous studies on pH_2 solids grown from the gas phase in an enclosed cell near 10 K

$\Rightarrow \quad \bar{\kappa} \approx 4000$ mW/cm-K

& $\quad L_{ph} \sim 0.1$ cm.

$$\kappa = C\, v\, L_{ph} / 3$$

[V.G. Manzhelii, B.Ya. Gorodilov, and A.I. Krivchikov, Low Temp. Phys. **22**, 131 (1996)]

Suggests $L_{ph} \sim 1$ μm in our rapid vapor deposited solids.

Summary

* Demonstrated production of millimeters-thick transparent pH_2 solids by rapid vapor deposition.

* Demonstrated that vapor deposited pH_2 solids are densest close-packed solids, NOT amorphous.

* Demonstrated suitability of vapor deposited pH_2 solids as hosts for high resolution IR absorption spectroscopy of chemically interesting dopants.

* Generalized phenomenon of dopant induced IR activity.

* Exploited CO/pH_2 spectroscopy to probe sample temperature during deposition, and to estimate thermal conductivity.

Collaborators

* Mr. Simon Tam and Ms. Michelle E. DeRose, AFRL/PRSP
  responsible for our experimental data.

* Prof. Takamasa Momose, Kyoto U.
  spectroscopy of CH_4, C_{60}, and CO doped pH_2 solids.

* Prof. Robert J. Hinde, U. Tennessee at Knoxville
  dopant-induced IR absorptions.

* Prof. David T. Anderson, U. Wyoming
  spectroscopy of $(HCl)_2$ in solid pH_2.